


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




3895
3896
3897
3898
3899
3900
3901
3902
3903
3904
3905
3906
3907
3908
3909
3910
3911
3912
3913

Shoib Ahmad Malik Baba. Phytochemical analysis and antioxidant activity of different tissue types of <i>Crocus sativus</i> and oxidative stress alleviating potential of saffron extract in plants, bacteria, and yeast. <i>South African Journal of Botany</i> . 2015. 99:80-87
G. M. Yang Bridger. Cold Acclimation Increases Tolerance of Activated Oxygen in Winter Cereals. <i>Journal of Plant Physiology</i> . 1994. 144:235-240
Kunmei Li Chen. TaSCL14, a Novel Wheat (<i>Triticum aestivum</i> L.) GRAS Gene, Regulates Plant Growth, Photosynthesis, Tolerance to Photooxidative Stress, and Senescence. <i>Journal of Genetics and Genomics</i> . 2015. 42:21-32
Ill-Min Park Chung. Resveratrol accumulation and resveratrol synthase gene expression in response to abiotic stresses and hormones in peanut plants. <i>Plant Science</i> . 2003. 164:103-109
Carolina Culiáñez-Macià Cortina. Tomato abiotic stress enhanced tolerance by trehalose biosynthesis. <i>Plant Science</i> . 2005. 169:75-82
Yong-Sheng Kong Deng. Heterology expression of the tomato LeLhcb2 gene confers elevated tolerance to chilling stress in transgenic tobacco. <i>Plant Physiology and Biochemistry</i> . 2014. 80:318-327
Angela Fletcher Gilley. Gibberellin antagonizes paclobutrazol-induced stress protection in wheat seedlings. <i>Journal of Plant Physiology</i> . 1998. 153:200-207
Upadhyaya Hemavathi. Over-expression of strawberry d-galacturonic acid reductase in potato leads to accumulation of vitamin C with enhanced abiotic stress tolerance. <i>Plant Science</i> . 2009. 177:659-667
Ěva Inaba Hideg. Dark adapted leaves of paraquat-resistant tobacco plants emit less ultraweak light than susceptible ones. <i>Biochemical and Biophysical Research Communications</i> . 1991. 178:438-443
Karen W. Negrotto Hughes. Free-radical stress response in paraquat-sensitive and resistant tobacco plants. <i>Environmental and Experimental Botany</i> . 1984. 24:151-157
Kohtaro Homma Iseki. The effects of cross-tolerance to oxidative stress and drought stress on rice dry matter production under aerobic conditions. <i>Field Crops Research</i> . 2014. 163:18-23
Pavanasam Rajamanickam Nithya Meenakshi Jeyaramraja. RETRACTED: Water deficit induced oxidative damage in tea (<i>Camellia sinensis</i>) plants. <i>Journal of Plant Physiology</i> . 2005. 162:413-419
Amr R. A. Schei Kataya. MAP kinase phosphatase 1 harbors a novel PTS1 and is targeted to peroxisomes following stress treatments. <i>Journal of Plant Physiology</i> . 2015. 179:12-20
Takao Matsui Koeduka. Rice fatty acid -dioxygenase is induced by pathogen attack and heavy metal stress: activation through jasmonate signaling. <i>Journal of Plant Physiology</i> . 2005. 162:912-920
Boo-Ja Kim Lee. Pathogen-inducible CaUGT1 is involved in resistance response against TMV infection by controlling salicylic acid accumulation. <i>FEBS Letters</i> . 2009. 583:2315-2320
Huizhi Wang Liu. A rice serine carboxypeptidase-like gene OsBISPL1 is involved in regulation of defense responses against biotic and oxidative stress. <i>Gene</i> . 2008. 420:57-65
Jia Shi Liu. Copper-induced hydrogen peroxide upregulation of a metallothionein gene, OsMT2c, from <i>Oryza sativa</i> L. confers copper tolerance in <i>Arabidopsis thaliana</i> . <i>Journal of Hazardous Materials</i> . 2015. 294:99-108
Wondi Norman Mersie. Response of beans and hairy fleabane leaves to ozone and paraquat with and without the antiozonant, ethylenediurea. <i>Environmental and Experimental Botany</i> . 1994. 34:379-383
Shigeto Nakatani Morita. Differential Expression of Two Cytosolic Ascorbate Peroxidases and Two Superoxide Dismutase Genes in Response to Abiotic Stress in Rice. <i>Rice Science</i> . 2011. 18:157-166

Crocus sativus L. belongs to family Iridaceae and its stigma part which forms commercial saffron has been used as a spice
Summary Injury from freezing stress may be caused by degradative reactions initiated by activated oxygen. The relationsh
Rates of photosynthesis, tolerance to photooxidative stress, and senescence are all important physiological factors that a
The peanut is one of the limited number of plant species that synthesize resveratrol, which is both a phytoalexin with ant
Trehalose is a nonreducing disaccharide of glucose that has been correlated with tolerance to different stress conditions.
Chilling is one of the most serious environmental stresses that disrupt the metabolic balance of cells and enhance the pro
Summary Gibberellic acid (GA) antagonism of the growth inhibitory and stress protective effects of paclobutrazol (P) was
Vitamin C (ascorbic acid) is an essential component for collagen biosynthesis and also for the proper functioning of the c
Long term light emission was compared from leaves of paraquat-resistant and -susceptible tobacco plants. In the minute
Plants regenerated from paraquat-resistant tobacco cell lines and their sexual progeny were evaluated for paraquat resis
Oxidative damage occurring in plant cells under drought stress is a known cause of reduced plant primary production. De
This article has been retracted at the request of the chief editors and author. Reason: this article contains material that h
In Arabidopsis thaliana, twenty mitogen-activated protein kinases (MAPKs/MPKs) are regulated by five MAP kinase phosph
Summary Plant fatty acid α -dioxxygenases (DOXs) catalyze the stereospecific conversion of fatty acids into the correspond
Capsicum annuum L. Bugang exhibits a hypersensitive response against Tobacco mosaic virus (TMV) P0 infection. The C.
Serine carboxypeptidase-like proteins (SCPLs) comprise a large family of protein hydrolyzing enzymes that play roles in m
Metallothioneins (MTs) are low-molecular-weight, cysteine-rich metal-binding proteins found in numerous genera and s
Growth chamber experiments were conducted to determine if there is a pattern of cross-tolerance to paraquat and ozon
Superoxide dismutase (SOD) and ascorbate peroxidase (APX) play central roles in the pathway for scavenging reactive ox

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



3914
3915
3916
3917
3918
3919
3920
3921
3922
3923
3924
3925
3926
3927
3928
3929
3930
3931
3932
3933

Hee-Yeon Seok Park. AtERF71/HRE2 transcription factor mediates osmotic stress response as well as hypoxia response in Arabidopsis. <i>Biochemical and Biophysical Research Communications</i> . 2011. 414:135-141
Douglas E. Heath Ryerson. Fungal elicitation of wall modifications in leaves of <i>Phaseolus vulgaris</i> L. cv. Pinto I. Attempts to mimic elicitation with chemical treatments. <i>Physiological and Molecular Plant Pathology</i> . 1992. 40:271-281
J. S. Kim Shin. Resistance levels and fitness of glufosinate-resistant transgenic sweet potato in field experiments. <i>Field Crops Research</i> . 2011. 121:324-332
De-Feng Wang Shu. Antisense-mediated depletion of tomato chloroplast glutathione reductase enhances susceptibility to chilling stress. <i>Plant Physiology and Biochemistry</i> . 2011. 49:1228-1237
C. J. Preston Soar. Polyamines can inhibit paraquat toxicity and translocation in the broadleaf weed <i>Arctotheca calendula</i> . <i>Pesticide Biochemistry and Physiology</i> . 2004. 80:94-105
Nakarin Kumla Suwannarach. Characterization and efficacy of <i>Muscodor cinnamomi</i> in promoting plant growth and controlling <i>Rhizoctonia</i> root rot in tomatoes. <i>Biological Control</i> . 2015. 90:25-33
Zoltán Darkó Szigeti. Diquat Resistance of Different Paraquat Resistant <i>Conyza canadensis</i> (L.) Cronq. Biotypes. <i>Journal of Plant Physiology</i> . 1994. 144:686-690
Gyula Darkó Váradi. Xanthophyll Cycle Patterns and in vivo Photoinhibition in Herbicide-Resistant Biotypes of <i>Conyza canadensis</i> . <i>Journal of Plant Physiology</i> . 1994. 144:669-674
Li-Yan Li Wang. Antisense-mediated depletion of tomato GDP-l-galactose phosphorylase increases susceptibility to chilling stress. <i>Journal of Plant Physiology</i> . 2013. 170:303-314
Deepanker Ahmed Yadav. Genome-wide identification and expression profiling of annexins in <i>Brassica rapa</i> and their phylogenetic sequence comparison with <i>B. juncea</i> and <i>A. thaliana</i> annexins. <i>Plant Gene</i> . 2015. 4:109-124
Jingli Niu Yan. Arabidopsis CIPK14 positively regulates glucose response. <i>Biochemical and Biophysical Research Communications</i> . 2014. 450:1679-1683
Qin Huang Yu. Direct measurement of paraquat in leaf protoplasts indicates vacuolar paraquat sequestration as a resistance mechanism in <i>Lolium rigidum</i> . <i>Pesticide Biochemistry and Physiology</i> . 2010. 98:104-109
Maoying Pan Zhang. ZmMKK3, a novel maize group B mitogen-activated protein kinase kinase gene, mediates osmotic stress and ABA signal responses. <i>Journal of Plant Physiology</i> . 2012. 169:1501-1510
P. G. Luyen Potvin. Electrostatic bubbles and supramolecular assistance of photosensitization by carboxylated Ru(II) complexes. <i>J Am Chem Soc</i> . 2003. 125:4894-906
L. Talarico Flamigni. Photoinduced electron transfer between the interlocked components of porphyrin catenanes: effect of the presence of nonequivalent reduction sites on the charge recombination rate. <i>Chemistry</i> . 2003. 9:2649-59
L. Bouanda Dupont. Metal ions binding onto a lignocellulosic substrate extracted from wheat bran: a NICA-Donnan approach. <i>J Colloid Interface Sci</i> . 2003. 263:35-41
V. D. Lagunova Samuilov. Role of chloroplast photosystems II and I in apoptosis of pea guard cells. <i>Biochemistry (Mosc)</i> . 2003. 68:912-7
C. Tabares Bittel. The oxidant-responsive diaphorase of <i>Rhodobacter capsulatus</i> is a ferredoxin (flavodoxin)-NADP(H) reductase. <i>FEBS Lett</i> . 2003. 553:408-12
V. M. Mattammal Lakshmi. Mechanism of peroxidative activation of the bladder carcinogen 2-amino-4-(5-nitro-2-furyl)-thiazole (ANFT): comparison with benzidine. <i>Carcinogenesis</i> . 1990. 11:1965-70
H. R. Vignon Tseng. Redox-controllable amphiphilic [2]rotaxanes. <i>Chemistry</i> . 2004. 10:155-72

Various transcription factors are involved in the response to environmental stresses in plants. In this study, we character
Based on results from previously published work, various chemical solutions were injected into the intercellular spaces o
Transgenic herbicide-resistant sweet potato plants [<i>Ipomoea batatas</i> (L.) Lam.] produced through a biolistic transformati
A tomato (<i>Lycopersicon esculentum</i> Mill.) chloroplast glutathione reductase gene (LeGR) was isolated and antisense tran
Reduced paraquat transport from the site of application to the site of action in the chloroplast seems a likely mechanism
<i>Muscodor cinnamomi</i> was selected and investigated for its in vitro ability to produce indole-3-acetic acid (IAA) to solubiliz
Summary Photosynthetic responses of paraquat/atrazine coresistant (PqAR) and only paraquat resistant (PqR) biotypes c
Summary The xanthophyll cycle and in vivo photoinhibition were investigated in the herbicide-susceptible (S), paraquat-r
The GDP-l-galactose phosphorylase (GGP), which converts GDP-l-galactose to l-Gal-1-phosphate, is generally considered t
Abiotic stresses affect the yield of crop plants worldwide. Plant species have evolved in such a way that they are able to c
Calcium is a ubiquitous intracellular secondary messenger in plants. Calcineurin B-like proteins (CBLs), which contain fou
Sequestration of paraquat away from its target site in the chloroplast has been proposed as a mechanism of paraquat res
Mitogen-activated protein kinase (MAPK) cascades are important intracellular signaling modules and function as a conve
The paper examines the supramolecular effects at play during photosensitization by carboxylated Ru(II) sensitizers, both
[2]Catenanes made up of several polyether-strapped porphyrin macrocycles interlinked with the cyclic electron acceptor
In this study ion binding to solid organic matter was investigated. We used the NICA-Donnan model to describe the inter
Received Revision received We investigated the CN--induced apoptosis of guard cells in epidermal peels isolated from pe
Challenge of <i>Rhodobacter capsulatus</i> cells with the superoxide propagator methyl viologen resulted in the induction of a
The mechanism of activation of the bladder carcinogen 2-amino-4-(5-nitro-2-furyl)thiazole (ANFT) was investigated by co
With the fabrication of molecular electronic devices (MEDs) and the construction of nanoelectromechanical systems (NE

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant